

**Table 4.5-8 STOMACH CLUSTER ANALYSIS –Dominant Prey Species Responsible for Clustering Structure**

Cluster	1								2						3			outlier	
	a				b				c	a				b	c		a	b	
Location/Lane	LT3	LT3	MH3	MH1	LT1	MH1	MH3	MH3	LT1	MH3	MH3	MH1	MH1	LT1	LT1	LT1	LT1	LT3	MH3
Tow Fish	6	3	6	1	2	6	6	1	1	3	3	1	3	1	2	6	6	1	1
	YT	YT	YT	YT	YT	YT	WF	WF	YT	YT	WF	WF	WF	WF	WF	WF	YT	WF	YT
Spionidae	100	100	100	100	100	100	100	100	30	30	30	30	5	5	1	1	30	1	5
Maldanidae	5	5	5	30	5	5	30	100	5	5	30	30	5	5	5	5	5	1	1
Caprellidae		5	5	5	30	30	30	5		5	30	30	30	5	5	100	100	30	5
Ampharetidae	30	5	5	5	30	30	30	30	5	5	5	5	5	5	30	30	30	100	1
Aoridae ( <i>Unciola</i> )	30	30	5	1	30	30	1	1	5	1		5	5	5	5	5	30	5	1
Lumbrineridae	1	5	5	5	5	1	30	5	5	1	5	5	5	5	30	5	30	5	